

## Coprophilic dipteran community associated with horse dung in Malaysia

\*Heo C.C.<sup>1,2</sup>, Latif, B.<sup>1</sup>, Kurahashi H.<sup>3</sup>, Hayashi T.<sup>4</sup>, Nazni W.A.<sup>5</sup> and Omar B.<sup>6</sup>

<sup>1</sup>Faculty of Medicine, Universiti Teknologi MARA, Sungai Buluh Campus, Jalan Hospital, 47000 Sungai Buluh, Selangor, Malaysia

<sup>2</sup>Institute of Forensic Science (IFS), Universiti Teknologi MARA, Level 4, Academic Building, Faculty of Medicine, 47000 Sungai Buluh, Selangor, Malaysia

<sup>3</sup>International Department of Dipterology (IDD), Hikawadai 1-2-21, Higashikurume City, Tokyo 203-0004, Japan

<sup>4</sup>Department of Medical Entomology, National Institute of Infectious Diseases, Toyama 1-23-1, Shinjuku-Ku, Tokyo 162-8640 Japan

<sup>5</sup>Medical Entomology Unit, Institute for Medical Research (IMR), Jalan Pahang, 50588 Kuala Lumpur, Malaysia

<sup>6</sup>Department of Biomedical Sciences, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Jalan Raja Muda Abdul Aziz, 50300 Kuala Lumpur, Malaysia

(email: \*chin@salam.uitm.edu.my)

### Abstract

Blood-sucking flies are known as vectors of pathogens to farm animals worldwide. However, little is known about dipteran biodiversity associated with horse dung in Malaysia which could serve as vectors. Thus, a field trip to a horse farm located in Tanjung Rambutan, Perak, peninsular Malaysia was conducted in 2010. We examined adults and larvae of dipterans associated with 100 horse dung pats. A total of 1480 dipteran specimens from nine families were collected including Sphaeroceridae, Muscidae, Sarcophagidae, Calliphoridae, Sepsidae, Tabanidae, Ulidiidae, Dolichopodidae and Milichiidae. The lesser dung fly (Sphaeroceridae) which consisted of four species was the most abundant dipteran followed by sarcophagids and muscids. Seven species of muscids were collected from the horse farm and three of them were haematophagic namely *Musca conducens* Walker, 1859, *Musca ventrosa* Wiedemann, 1830 and *Stomoxys calcitrans* (Linnaeus, 1758). Fly larvae collected from the dung were raised to adult stage and subsequently identified as *M. conducens* and *Neomyia gavis* Walker, 1859. We reported new locality records for three species of Muscidae in Malaysia namely *Pyrellia proferens* Walker, 1859, *Lispe kowarzi* Becker, 1903 (new to peninsular) and *N. gavis* Walker, 1859.

**Keywords:** Coprophilic Diptera, horse dung, Malaysia, new record, *Pyrellia proferens*, *Lispe kowarzi*, *Neomyia gavis*.

Received: 24 February 2015; Revised: 22 April 2015; Online: 4 May 2015; Published: 5 November 2015.

### Introduction

The Diptera, or known as two-winged flies, is one of the largest orders of Insecta and can be found almost ubiquitously on Earth (Triplehorn and Johnson, 2005). Members of Diptera play variety of roles in ecosystem either as pollinators, nutrient recyclers, or carrion decomposers (Borkent and Harder, 2007; Hirabayashi and Wotton, 1998; Putman, 1978). However, there are many dipteran species of economic importance which are responsible for loss of agricultural products and commodity (White and Elson-Harris, 1992; Spencer, 1973). Most importantly, flies have

been incriminated either as biological or mechanical vectors to wide range of infectious diseases that are prevalent in both developed and developing countries (Gubler, 2002).

Other than ticks (Acari: Ixodida), Diptera is considered a significant group of arthropods in term of medical and veterinary importance (Cleton *et al.*, 2012). Flies are found accountable for the transmission of variety of pathogens namely viruses, bacteria, protozoans and helminthes (Gestmann *et al.*, 2012), which are responsible for the spread of infectious diseases such as malaria, lymphatic

filariasis, dengue, yellow fever, West Nile fever, and chikungunya (transmitted by nematocerans such as mosquitoes), leishmaniasis, onchocerciasis, loiasis (transmitted by sand flies, black flies and deer flies, respectively) (Service, 2012). Other members of Diptera such as house flies (*Musca domestica* L.), face flies (*Musca autumnalis* De Geer), dog dung flies (*Musca sorbens* Wiedemann), and blow flies (*Lucilia* spp.) are a source of nuisance to humans and animals (Graczyk *et al.*, 2001). Some species may act as the causative agent to myiasis either obligatory or facultative on animal hosts (James, 1947; Hall and Wall, 1995). Besides, there are many dung-frequenting and carrion-breeding dipterans which may spread pathogenic organisms to the surrounding environment, including human dwelling (Howard, 2001).

Other than the diseases mentioned above, veterinary important Diptera have been reported to impose a serious economic loss to the agricultural sector (Bowman, 2014). For instance, In the United States, flies had impacted the cattle production substantially by losing \$2,211 million per year (Taylor *et al.*, 2012a). The stable fly, with just 200,000 flies emerging from an average sized winter hay feeding site could reduce annual milk production of 50 dairy cows by an estimated 890 kg (Taylor *et al.*, 2012b).

In Malaysia, Reid (1953) published the first note on the local house flies and blow flies and indicated their important roles as potential mechanical transmitter of pathogens causing human diseases. Subsequently, Sulaiman *et al.* (1988) demonstrated that the cyclorrhaphan flies collected from four sites in Malaysia are carriers of human parasitic helminths such as *Ascaris lumbricoides*, *Trichuris trichiura*, and filariform larva of hookworm (*Necator americanus*). Sulaiman *et al.* (2000) trapped four species of synanthropic flies in downtown Kuala Lumpur viz. *Chrysomya megacephala*, *Chrysomya rufifacies* (= *Achoetandrus rufifacies*), *M. domestica* and *M. sorbens*. A total of 18 bacterial species was isolated from these specimens, with *Burkholderia pseudomallei*, the causative pathogen for melioidosis, had been reported for the first time. Nazni *et al.* (2005) examined bacteria from the house flies and found 10 different species of bacteria (e.g., *Bacillus* sp., *Staphylococcus* sp., *Proteus* sp.,

*Escherichia* sp., *Klebsiella* sp. etc.) isolated from feces, vomitus, external surfaces and internal organs of house flies. However, newly emerged house fly did not harbor any bacteria. The importance of house fly wings in mechanical transmission of *Vibrio cholerae* was assessed and the results revealed that the wings did not play an important role in pathogen transmission (Yap *et al.*, 2008). Nazni *et al.* (2013) carried out a study on house fly to establish whether the house flies can transmit the H1N1 virus mechanically. The findings indicated that the persistency of H1N1 virus on fly legs could be detected up to 24 hours either in chilled or actively flying flies. However, virus was not found in the vomitus, feces and the external body of the flies.

Recently, filth fly surveys were conducted in Malaysia by Nurita and Hassan (2013), Khoso *et al.* (2015) and Heo *et al.* (2010) where they found a very high diversity of filth flies in the region. Nurita and Hassan (2013) recorded eight dipteran species breeding in the solid waste namely: *M. domestica*, *M. sorbens*, *Synthesiomia nudisetia* Van der Wulp, *Hydrotaea chalcogaster* (Wiedemann), *C. megacephala*, *Lucilia cuprina* (Wiedemann), *Hemipyrellia ligurriens* (Wiedemann) and *Sarcophaga* sp.; whereas Heo *et al.* (2010) recorded 16 families of Diptera from the cow dung. These results showed that landfills and animal farms can act as major breeding grounds for filth flies.

From the above stated data, it is clear that little is known about the coprophilic dipteran community associated with horse manure in Malaysia, particularly the blood sucking flies that might feed on horses. Hence, there is a need to identify the veterinary important Diptera in the horse farm, as the baseline data would be of paramount important in terms of risk management and in vector control programs in future. The present study was aimed to determine the diversity and abundance of coprophilic Diptera associated with horse manure, and to identify the veterinary and medically important dipterans in a horse farm in Malaysia.

## Materials and Methods

Horse farm located in Tanjung Rambutan, Perak, Malaysia was selected as the study site (4°41'13" N 101°09'31" E, ~89 m a.s.l.) located approximately 212 km northwest

from the capital city, Kuala Lumpur (Fig. 1). A total of four visits to the horse farm were conducted from 8-11 November 2010. Duration and time of fly collection was 5 hours per visit from 9 am to 2 pm. The mean temperature and humidity during the survey period was  $28.92 \pm 2.36$  °C and  $73.88 \pm 10.72\%$ , respectively. The weather conditions during the four-day field trials were mostly cloudy and all weather data was obtained from the nearest weather station in Ipoh, Perak.

We examined a total of 100 horse dung pats scattered in an area of approximately 5,900 m<sup>2</sup>. The age of horse dung examined was from fresh to 3-day old. Each dung pat was examined for the presence of adult flies and dung-breeding larvae. Adult flies either resting on or swarming around the dung pats were collected using a sweep net or transparent plastic bags, which were then transferred to a killing jar which contained cotton balls soaked with ethyl acetate. A pair of forceps was used to examine and recover fly larvae from the dung pats. Any larvae or soil arthropods observed in the dung were then collected and preserved in vials (diameter: 25 mm; height: 85 mm) containing 70% ethanol with label. Several larvae from the same batch were also collected and placed in a transparent plastic container (diameter: 44 mm; height: 57 mm; volume: 70 ml) together with a small amount of horse manure which served as food source for rearing purposes. The screw cap of the rearing container was removed and replaced with a breathable paper towel which had been tightened with a rubber band around the lid. Few drops of water were added *ad-libitum* to maintain the moisture of the horse manure throughout the rearing periods. All preserved and rearing specimens were then transferred to parasitology laboratory at the Faculty of Medicine, Sungai Buluh Campus, Universiti Teknologi MARA.

The emerging adult flies from rearing containers were prompted for species identification under a stereomicroscope (Olympus SZX7, Japan) by using the keys provided in Triplehorn and Johnson (2005), Emden (1965) and Kurahashi *et al.* (1997). Several adult specimens from the families Calliphoridae, Muscidae and Sarcophagidae were sent to the third author (Kurahashi, H.) while specimens of Sphaeroceridae were sent to the fourth author (Hayashi, T.) for species confirmation. Other soil arthropods recovered

from horse dung were processed for preservation and identified by using respective taxonomical keys to the lowest taxon. Collembola, ants and centipedes were preserved in vials filled with 70% ethanol while Coleoptera were pinned and labelled. For acari processing, the mite specimens were placed in lactophenol (50% lactic acid, 25% phenol crystal, and 25% distilled water) for clearing purposes. They were left in clearing medium for a week and then transferred on slides by using a probe. Hoyer's medium was used as the mounting fluid and then the specimens were covered with thin cover slips. The slides were then placed in a drying oven at 40°C for a week. The mites were identified to family level based on Krantz and Walter (2009) whereas genus of Macrochelidae was confirmed using the key of Emberson (1980).

Diversity and abundance data of the collected dipterans was calculated for ecological indices (Dominance, Species Richness, Simpson's Index, Shannon-Wiener Index, and Evenness) for the study period. Dominance ( $D_i$ ) was calculated according to the equation:

$$D_i = \frac{n_i}{N} \times 100$$

Where  $n_i$  is the number of individuals collected during the study period, and  $N$  is the total number of specimen collected. Species dominance of all families or species were classified according to Tischler's scale: eudominant  $10\% \leq D_i \leq 100\%$ , dominant  $5\% \leq D_i \leq 10\%$ , subdominant  $2\% \leq D_i \leq 5\%$ , recedent  $1\% \leq D_i \leq 2\%$ , and subrecedent  $0\% \leq D_i \leq 1\%$  (Tischler, 1949).

Species richness ( $S$ ) is the total of different species presented in the sample while Simpson's Index ( $D$ ) measured both richness and proportion of each species and is calculated by using the formula:

$$D = \sum_{i=1}^S P_i^2$$

Where  $P_i$  is the proportion of species  $i$ . In brief, Simpson's index is the sum of proportion of each species in the community and represented the probability of two randomly selected individuals in the community belong to the same species. Shannon-Wiener Index ( $H'$ ) is

similar with Simpson's Index where the measurement takes species richness and proportion of species into account, and is calculated by the following formula:


$$H' = - \sum_{i=0}^n P_i (\ln P_i)$$

In general, Shannon-Wiener Index is the negative sum of multiply products between species proportion ( $P_i$ ) and natural log of species proportion ( $\ln P_i$ ). Evenness (E) is an

indicator of similarity in abundance of different species. Evenness is measured on the scale from 0 to 1 where zero represents more variation in communities whereas one represents complete evenness. Evenness is defined as:

$$E = \frac{H'}{\ln S}$$

Evenness is the number obtained via dividing the value of Shannon-Wiener Index by natural log of species richness (S).



**Fig. 1. Location of the study site at peninsular Malaysia is indicated by a pink arrow (map created using www.mapbox.com)**

## Results

A total of 1,480 adult specimens from nine families of Diptera were collected from the horse farm. The families consisted of Sphaoceridae (four species), Muscidae (seven species), Sarcophagidae (two species), Calliphoridae (two species), Sepsidae (two species), Tabanidae (one species), Ulidiidae (one species), Dolichopodidae (one species) and Milichiidae (one species) (Table 1). Throughout the study period, the most abundant dipterans caught were Sphaeroceridae (93.78%), followed by Sarcophagidae (3.04%), Muscidae (2.30%), Calliphoridae (0.50%), Sepsidae and Ulidiidae (0.10% each) and the least were Tabanidae, Dolichopodidae and Milichiidae (0.06% each) (Fig. 2).

We collected 34 individual of muscids which consisted of seven species (Fig. 3). The species of muscid flies according to their decreasing abundance were as follows: *M. domestica* (32.4%), *Musca conducens* Walker (26.50%), *Lispe kowarzi* Becker (23.50%), *Pyrellia proferens* Walker (5.90%), *Neomyia gavis* Walker (5.90%), *Musca ventrosa* Wiedemann (2.90%) and *Stomoxys calcitrans* (2.90%). Out of seven species recovered, three species were blood sucking muscids (*M. conducens*, *M. ventrosa* and *S. calcitrans*), with *M. conducens* being the most abundant haematophagic species in the horse farm.

Three species of Muscidae were also recorded for the first time in Malaysia namely *L. kowarzi* (new in peninsular Malaysia), *P.*

## Coprophilic dipteran community associated with horse dung in Malaysia

*proferens*, and *N. gavis*a (Fig. 4). These adult flies were collected on or around the horse dung pats, indicating their coprophilic behavior in nature. We hereby provide locality notes for all three species:

### Specimens examined:

*Lispe kowarzi* Becker, 1903

MALAYSIA: Perak: Tanjung Rambutan 4°41'13" N 101°09'31"E, ~89 m a.s.l. 9.xi.2010, C.C. Heo, 4 males, 4 females.

*Pyrellia proferens* (Walker, 1859)

MALAYSIA: Perak: Tanjung Rambutan 4°41'13" N 101°09'31"E, ~89 m a.s.l. 10.xi.2010, C.C. Heo, 1 male, 1 female.

*Neomyia gavis*a (Walker, 1859)

MALAYSIA: Perak: Tanjung Rambutan 4°41'13" N, 101°09'31"E, ~89 m a.s.l. 10.xi.2010, C.C. Heo, 1 female.

The most abundant Diptera caught in this study was the lesser dung flies (Sphaeroceridae), which constituted about 94% of total specimen collected. This family was observed to frequent on almost every horse dung scattered in the farm, and usually seen in a large group resting on dung surfaces. Four species of sphaerocerids were collected and identified: *Norrbomia tropica* Duda, *Coproica rufifrons* Hayashi, *Coproica coreana* Papp and *Coproica aliena* Papp (Fig. 5). On the other hand, two species of flesh flies (Sarcophagidae) were found in the farm namely *Parasarcophaga taenionota* (Wiedemann) and *Liopygia ruficornis* (Fabricius), and two species of blow flies (Calliphoridae) were identified associated with horse manure, the oriental latrine fly, *Chrysomya megacephala* Fabricius and the hairy maggot blow fly, *Achoetandrus rufifacies* (Macquart). These are common blow fly species found in Malaysia and were known to visit decomposing organic matter. Other than that, we also collected specimens from the families Sepsidae, Tabanidae, Ulidiidae (*Physiphora* sp.), Dolichopodidae and Milichiidae (*Milichiella* sp.). Note that the tabanid fly is also a blood-sucking species, however, it was rarely seen in the farm.

In addition to Diptera, there were many other taxa associated with horse feces. Examination into the dung pats revealed

springtails (one family), beetles (four families), mites (three families), centipedes (one order), ants and spider (one family). However, no earthworm was collected. A butterfly (Lepidoptera: Lycanidae) was also observed to frequent on dung surface. Table 2 listed the families of non-Diptera taxa associated with horse dungs.

Results of ecological indices obtained from formulas stated in methodology were presented in Table 3 and 4. Among dipteran families, family Sphaeroceridae was the most dominant family. Simpson's Index was 0.88 while Shannon-Wiener Index was 0.31. The abundance of all the families collected was dissimilar and skewed, resulted in an uneven distribution among families ( $E= 0.14$ ). In terms of Family level, only Muscidae was analyzed. *Musca domestica*, *M. conducens* and *L. kowarzi* were all classified in the eudominant group. Simpson's Index and Shannon-Wiener Index were 0.26 and 1.54, respectively, indicated a more equal proportion among species. This observation was strengthened with a higher evenness index ( $E=0.80$ ).

### Discussion

Animal dung pats are patchy but resourceful ephemeral microhabitats which host a vast diversity of species (Hanski and Koskela, 1977). The adults and larvae of Coleoptera and Diptera are considered to play a major role in the dung utilization. However, little is known about the coprophilous fly species compared to studies conducted on dung beetles (Hammer, 1941; Nichols *et al.*, 2007).

Sphaeroceridae are very small, black or brown flies that can be identified by the swollen hind tarsi. They can be found in humid and swampy places near excrement and often aggregated in large number on various mammalian dung piles (Laurence, 1955; Triplehorn and Johnson, 2005). Unexceptionally, the most abundant Diptera collected on horse dung in this study was sphaerocerids. Similar observation was made in U.K. on stable manure and two species of sphaerocerids were considered as potential pests (Hussey, 1957). This finding was in agreement with Bai and Sankaran (1977) where sphaerocerids were found breeding in bovine manure in India. Similar results were demonstrated in North America where

sphaerocerids were collected from cattle droppings (Blume, 1985). Other than cattle and horse manures, sphaerocerids can be found in poultry manure (Hulley, 1986). Sphaerocerids, together with acarid mites were also reported as the first colonizer on poultry manure (Stoffolano and Geden, 1987). The cosmopolitan genus *Coproica* is known to be endemic on various kinds of pasture dung whereas some species breed in decaying

vegetable material. In fact, most larvae actually feed on microbial layer developing on the decaying matter (Papp, 2008). However, during the present study, we were not able to collect any sphaerocerid larva in the horse manure. Perhaps it was due to the smaller sample size or other abiotic factors (e.g., condition of horse manure or dung management strategies employed by the farm).

**Table 1. Family and species of Diptera recovered from horse dung**

Family	Species
Sphaeroceridae	<i>Norrbomia tropica</i> Duda, 1923 <i>Coproica rufifrons</i> Hayashi, 1991 <i>Coproica coreana</i> Papp, 1979 <i>Coproica aliena</i> Papp, 2008
Muscidae	<i>Musca domestica vicina</i> Macquart, 1850 <i>Musca conducens</i> Walker, 1859 <i>Musca ventrosa</i> Wiedemann, 1830 <i>Stomoxys calcitrans</i> (Linnaeus, 1758) <i>Pyrellia proferens</i> Walker, 1859 <i>Lispe kowarzi</i> Becker, 1903 <i>Neomyia gavis</i> Walker, 1859
Sarcophagidae	<i>Parasarcophaga taenionota</i> (Wiedemann, 1819) <i>Liopygia ruficornis</i> (Fabricius, 1794)
Calliphoridae	<i>Chrysomya megacephala</i> Fabricius, 1794 <i>Achoetandrus rufifacies</i> (Macquart, 1842)
Sepsidae	Unidentified sp. 1 Unidentified sp. 2
Tabanidae	Unidentified sp.
Ulidiidae	<i>Physiphora</i> sp.
Dolichopodidae	Unidentified sp.
Milichiidae	<i>Milichiella</i> sp.

**Table 2. Other arthropods associated with horse dungs**

Class	Order	Family	Subfamily / Species
Insecta	Collembola	Hypogastruridae	<i>Ceratophysella</i> sp.
	Coleoptera	Scarabaeidae	<i>Onthophagus</i> sp. Aphodiinae
		Hydrophilidae	Unidentified sp.
		Histeridae	Unidentified sp.
Staphylinidae		Staphylininae	
	Hymenoptera	Formicidae	<i>Odontoponera</i> sp.
Arachnida	Mesostigmata	Macrochelidae	<i>Macrocheles</i> sp.
		Uropodidae	Unidentified sp.
		Parasitidae	Unidentified sp.
	Araneae	Lycosidae	Unidentified sp.
Chilopoda	Geophilomorpha	Unidentified	Unidentified sp.

**Coprophilic dipteran community associated with horse dung in Malaysia**

**Table 3. Standard ecological indicators of Order Diptera found on horse dung**

Family	Number	Dominance	Tischler's scale	$P_i$	$P_i^2$	$P_i \ln(P_i)$
Sphaeroceridae	1387	93.7	Eudominant	0.937	0.878	-0.061
Sarcophagidae	45	3.0	Subdominant	0.030	0.001	-0.106
Muscidae	34	2.3	Subdominant	0.023	0.001	-0.087
Calliphoridae	7	0.5	Subrecedent	0.005	0.000	-0.025
Sepsidae	2	0.1	Subrecedent	0.001	0.000	-0.009
Ulidiidae	2	0.1	Subrecedent	0.001	0.000	-0.009
Tabanidae	1	0.1	Subrecedent	0.001	0.000	-0.005
Dolichopodidae	1	0.1	Subrecedent	0.001	0.000	-0.005
Millichidae	1	0.1	Subrecedent	0.001	0.000	-0.005
Total	1480	100.0		1.000		
<b>Ecological Indicators</b>			<b>Value</b>			
Species Richness (S)			9			
Simpson Index (D)			0.880			
Shannon-Wiener Index (H')			0.312			
Evenness (E)			0.142			

**Table 4. Standard ecological indicators of Family Muscidae found on horse dung**

Species	Number	Dominance	Tischler's scale	$P_i$	$P_i^2$	$P_i \ln(P_i)$
<i>M. domestica</i>	12	35.3	Eudominant	0.382	0.146	-0.368
<i>M. conducens</i>	9	26.5	Eudominant	0.265	0.070	-0.352
<i>L. kowarzi</i>	8	23.5	Eudominant	0.235	0.055	-0.340
<i>P. proferens</i>	2	5.9	Dominant	0.059	0.003	-0.167
<i>N. gavisia</i>	1	2.9	Subdominant	0.029	0.001	-0.104
<i>M. ventrosa</i>	1	2.9	Subdominant	0.029	0.001	-0.104
<i>S. calcitrans</i>	1	2.9	Subdominant	0.029	0.001	-0.104
Total	34	100.0		1.000		
<b>Ecological Indicators</b>			<b>Value</b>			
Species Richness (S)			7			
Simpson Index (D)			0.256			
Shannon-Wiener Index (H')			1.538			
Evenness (E)			0.790			




Fig. 2. Abundance of Diptera collected according to family (in percentage)




Fig. 3. Species dominance of seven species of family Muscidae collected from cow dung






Fig. 4. Newly recorded Muscidae from peninsular Malaysia. A. *Pyrellia proferens*; B. *Lispe kowarzi*; C. *Neomyia gavis*




Fig. 5. Sphaeroceridae collected from horse dung. A. *Coproica aliena*; B. *Coproica coreana*; C. *Coproica rufifrons*; D. *Norrbomia tropica*. Scale bar = 0.5 mm for all figures

We collected four species of blood-sucking flies (three in Muscidae and one in Tabanidae) during the study period. Among these, *M. conducens* was the most abundant (26.5% of total muscid caught), followed by *M. ventrosa* and *S. calcitrans* (both were equally abundant), and lastly a *Tabanus* sp. As discussed earlier, the blood sucking Diptera are considered medical and veterinary important pests due to their potential in causing and transmitting vector-borne pathogens. *Musca conducens* was incriminated as the vector for stephanofilariasis, which had been reported to cause chronic eosinophilic dermatitis in scrotal area among Charolais bulls in France (Watrelet-Virieux and Pin, 2006) and transmit *Stephanofilaria assamensis* in India and Russia (Patnaik and Roy, 1969; Johnson *et al.*, 1981) and *Stephanofilaria kaeli* in peninsular Malaysia (Fadzil, 1973). Interestingly, we collected 13 larvae of *M. conducens* from horse manures and reared them to the adult stage, out of these four males and nine females emerged (male: female ratio 1:2).

*Musca ventrosa* was found in fresh markets in Malaysia, albeit its population was low (Khoso *et al.*, 2015). Nazni *et al.* (2007) conducted a study on distribution of diurnal and nocturnal dipterous flies in Putrajaya, Malaysia and collected six specimens of *M. ventrosa* out of 1,534 flies (0.04%) during the day. Despite feeding on wounds, sores and bites inflicted by other insects, these flies also frequent cow dung and their larvae breed in it (Patton, 1922). So far, little literature is available on the status of *M. ventrosa* as a vector for veterinary diseases although it is a haematophagous species (Sucharit and Tumrasvin, 1981). In our study, we collected a single adult *M. ventrosa*. Due to their low population, their feeding impact on horses seems to be limited.

Adults of *S. calcitrans* suck blood and inflict painful bites and thereby causing trouble to the confined livestock such as cattle and horses (Service, 2012). The fly causes interference with the normal feeding activity of cattle which could result in weight loss, decreased milk production, and possibly anemia. Moreover, stable flies have been found responsible for the mechanical transmission of several livestock diseases such

as equine infectious anemia and bovine anaplasmosis (Bay and Harris, 1988). Likewise, we collected only one adult *S. calcitrans*, which may suggest limited vectorial capacity due to their low population density.

Heo *et al.* (2010) conducted a study on cow dung Diptera in Kuala Lumpur, Malaysia and found *Musca inferior* Stein to be the most abundant (37%) blood-sucking muscid associated with cattle manure, followed by *S. calcitrans* (10.3%), *M. ventrosa* (9.4%), *M. crassirostris* (3.4%) and *Haematobia* sp. (0.8%). In comparison with the present study, the results showed that *M. ventrosa* and *S. stomoxys* were present in both study sites and were of intermediate abundance either on cattle or horse manures. However, we did not collect *M. inferior*, *M. crassirostris* and *Haematobia* from the horse dung. Additionally, the most abundant non-biting Diptera associated with cattle dung was Sepsidae, which was different in the current study. A comparison of dipteran composition with Heo *et al.* (2010) is presented in Table 5.

Although cattle and horses are herbivores, there are some remarkable differences in the diversity of arthropod assemblages on their wastes. The differences in results of arthropod communities between cattle and horse dungs could be due to two main reasons: biotic and abiotic factors. Biotic factors include the differences in microbiome in the animal dung (Dowd *et al.*, 2008; Girija *et al.*, 2013; Costa and Weese, 2012), which may affect directly the behavior of arthropods (Ezenwa *et al.*, 2012) or perhaps the subsequent colonization events (Shade *et al.*, 2013). The microbiome on the dung could be the direct result from the animal intestine itself or the contamination from the external environment (Girija *et al.*, 2013). Besides, arthropod visiting the dung could also deliver and contribute microbes to it (Estes *et al.*, 2013) which consequently initiate a change in arthropod successional sequence either due to the exploration of new resources (Pechal *et al.*, 2014; Finley *et al.*, 2015) or interkingdom signaling through volatile organic compounds released by bacteria (Tomberlin *et al.*, 2012). Moreover, ecological interactions such as mutualism, competition, predation and parasitism could shape the structures and

## Coprophilic dipteran community associated with horse dung in Malaysia

functions of the microbial and arthropod communities in the dung ecosystem (Valiela, 1974). The abiotic factors include the differences in spatial and temporal factors (e.g., geographical locations of the study sites and timing of study), weather factors (e.g., differences in temperature and relative humidity, event of rains), and most notably, is the difference in the physical structure and chemical composition between cattle and

horse manures. Cattle dungs had more moisture content (31%) than horse (27%) (Akhtar *et al.*, 2013). Besides, horse dungs were more alkaline (pH 9.4) compared to cattle (pH 5.2). The percentage of organic carbon, C:N ratio, total nitrogen and concentration of inorganic nitrogen were generally higher in cow dung in comparison with horse manure (Ajwa and Tabatabai, 1994).

**Table 5. Comparison of dipteran composition of cow and horse dungs.**

Result	Cow dung (modified from Heo <i>et al.</i> , 2010 with correction on the species of the Diptera larva collected)	Horse dung
Location of study	Sentul Timur, Kuala Lumpur (3°11' N 101°41' E)	Tanjung Rambutan, Perak (4°41'13" N 101°09'31" E)
Total family collected	16	9
Most abundant Diptera	Sepsidae (43.8%)	Sphaeroceridae (93.8%)
Total species collected in Muscidae	12	7
Total blood sucking species	6	4
Percentage of haematophagous muscid collected	51.5%	29.4%
Most dominant haematophagous muscid	<i>M. inferior</i>	<i>M. conducens</i>
Diptera larvae collected from dung	<i>M. inferior</i> Stein Psychodidae	<i>M. conducens</i> <i>N. gavis</i>

*Neomyia* spp. is common muscids to frequent on animal dung. Their larvae are found to be saprophagous, coprophagous (Couri *et al.*, 2006) and also serves as food source to many other carnivorous larvae living in the dung (Emden, 1965). The larvae of *Neomyia* and other dung fauna (e.g., dung beetles and earthworms) play a vital role in dung decomposition and nutrient recycling (Holter, 1977, 1979; Sommer *et al.*, 2001). In Malaysia, six species of *Neomyia* have been documented to date namely *Neomyia rufitarsi* (Stein), *Neomyia coerulea* Wiedemann, *Neomyia coeruleifrons* (Macquart), *Neomyia diffidens* (Walker), *Neomyia lauta* Wiedemann and *Neomyia indica* Robineau-Desvoidy (Emden, 1965; Heo *et al.*, 2010). With the finding of *N. gavis* during the present study, the number of species of *Neomyia* in Malaysia has increased to seven. *Neomyia gavis* was recorded in Sichuan Province, China as domi-

nant species and can be seen all year round (Wang and Feng, 2008). To date, no literature is available about the role of *Neomyia* in disease transmission.

*Pyrellia proferens*, a new record from Malaysia, was previously documented from India, Myanmar and Indonesia. The adults can be found on flowers and on decaying animal matters, especially on dungs. The larvae live in animal excrement and feed on carrion (Emden, 1965).

*Lispe* spp. are shore-living muscids and their larvae are aquatic in the environment (Shinonaga and Kano, 1989; Pont *et al.*, 2012). The adults *Lispe* are known to be highly predacious to small insects and potentially useful in biological control of black fly (Werner and Pont, 2006). We had a chance to collect *L. kowarzi* through this study and it is therefore newly recorded in peninsular Malaysia. It was previously recorded in Kota

Kinabalu, Sabah, Malaysian Borneo (Vikhrev, 2012a). In general, this species is widely spread in South Palaearctic and Oriental region (Tumrasvin and Shinonaga, 1982; Ebejer and Gatt, 1999; Bharti, 2008; Vikhrev, 2012a). It is interesting to note that this species could be the predator to other small insects associated with horse manure. So far, six species of *Lispe* (*Lispe pacifica* Shinonaga and Pont, *Lispe assimilis* Wiedemann, *Lispe pectinipes* Becker, *Lispe manicata* Wiedemann, *Lispe orientalis* Wiedemann) have been recorded in Malaysia, including the newly recorded *L. kowarzi* (Shinonaga and Pont, 1992; Nazni *et al.*, 2007; Kurahashi and Shinonaga, 2009; Chew *et al.*, 2012; Vikhrev, 2012b).

House flies, *M. domestica*, are responsible for the spread of infectious diseases such as typhoid, dysentery, diphtheria, leprosy, tuberculosis, intestinal parasitic infections in humans and are found to be mechanical vectors for various pathogenic bacteria, viruses and protozoans (Greenberg, 1973; Chaiwong *et al.*, 2014). Blow flies such as *C. megacephala* and *A. rufifacies* may act as facultative myiasis agents on the wounds of animals inflicted by injuries or insect bites (Sukontason *et al.*, 2005). Moreover, *C. megacephala* has been shown to have greater chances of finding helminth ova attached to their external surfaces compared to *M. domestica*. A study conducted in Ethiopia demonstrated that *A. rufifacies* acted as a vector of at least five helminthes parasites and four species of protozoan parasites (Getachew *et al.*, 2007). It is pertinent to mention here that these two species of blow flies are forensically important in Malaysia and other adjacent countries and their larvae are frequently encountered in forensic cases (Sukontason *et al.*, 2002; Lee *et al.*, 2004; Chen *et al.*, 2004; Wang *et al.*, 2008).

*Parasarcophaga taenionota* and *L. ruficornis* have been found to breed on human and animal carcasses in Thailand and Malaysia (Sukontason *et al.*, 2007; Tan *et al.*, 2010; Kumara *et al.*, 2012) and are used by forensic entomologists in determining the minimum post-mortem interval (mPMI) of a corpse. In medicine, the larvae of *L. ruficornis* have been documented to cause myiasis in the vagina of a comatose woman in Thailand (Sucharit *et al.*, 1981). Both *P. taenionota* and *L. ruficornis* had been recently collected at several fresh markets in Malaysia, indicating their

synantrophic behaviors that could contribute as potential mechanical vectors of pathogens (Khosro *et al.*, 2015). Moreover, Heo *et al.* (2010) collected *L. ruficornis* from cattle droppings, implying its coprophilous behavior.

Dung beetles (Coleoptera: Scarabaeidae) play a crucial role in dung decomposition and improve nutrient recycling and soil structure (Hanski and Cambefort, 1991). Both adults and larvae are dung feeders, and through the process of feeding, dung beetles facilitate a range of ecosystem services such as secondary seed dispersal, bioturbation, parasite suppression, plant growth enhancement, and soil fertilization (Nichols *et al.*, 2008). Higher diversity of dung beetles could accelerate nitrogen and carbon transfer from the grass-produced dung to the soil (Yoshihara and Sato, 2015). Predators such as rove beetles (Staphylinidae), hister beetles (Histeridae), centipedes, spiders and ants were also found in the horse manures. These predators were searching for fly eggs or insect larvae that were residing in the dung (Triplehorn and Johnson, 2005). Predatory staphylinids inflict heavy mortality on fly eggs, maggots, pupae and adults (Valiela, 1969). In tropical rain forest in Sarawak (Malaysian Borneo), Hanski (1983) revealed 60 Scarabaeinae and four Hybosorinae (necrophagous subfamily of Scarabaeidae), about 20 Hydrophilidae, six Histeridae and more than 150 Staphylinidae from dung and carrion. The water scavenging beetles (Hydrophilidae) are coprophagous, but their larvae are predators (Bøving and Henriksen, 1938).

Apart from Insecta, we also collected mites (Acari: Mesostigmata) from the family Macrochelidae, Parasitidae and Uropodidae, with macrochelids being the most abundant inhabitant of the horse dung. The result of the present study is similar to the earlier study by Axtell (1963) where Macrochelidae, Uropodidae, Parasitidae, Oribatidae and Laelapidae were collected from 211 samples of domestic animal manures (e.g. cattle, horses, sheep, chickens and ducks). Mesostigmata are free-living predators and known to predate on soil nematodes, Collembola and insect larvae (Koehler, 1999). Macrochelidae are active consumers of eggs and larvae of synantrophic Diptera and they might be useful as biological control for dung breeding flies (Krantz, 1983). Macrochelid mite can be found especially on

coprophagous Scarabaeidae (e.g., *Onthophagus* sp. and *Aphodius* sp.) (Glida and Bertrand, 2002). Both Parasitidae and Uropodidae are predacious on other soil arthropods and are closely associated with dung beetles through phoresy (Mašán and Halliday, 2009).

Numerous individuals of collembolans (Hypogastruridae: *Ceratophysella* sp.) were collected from the horse dung where they occurred in large aggregation. The members of Hypogastruridae feed on the decaying plant materials (Greenslade and Ireson, 1986). They may also be feeding on nematodes and other microscopic animals associated with decomposition (Chernova *et al.*, 2007). It should be noted that *Ceratophysella* spp. are beneficial as they contribute to the decomposition of organic matter (e.g., carrion and dung) and enhance nutrient recycling in agricultural land (Zhang *et al.*, 2012).

In conclusion, four species of blood sucking flies were identified in the horse farm in Malaysia with *M. conducens* being the most abundant haematophagic muscid. The most populous non-biting Diptera in the farm was the sphaerocerids. Proper management of animal manure is highly recommended to reduce population of blood sucking muscids and tabanids. Annual trapping and phenological studies of horse dung Diptera should be carried out to draw a clearer picture of the bionomics of veterinary important flies, as well as those dung decomposers who play a vital ecological role in dung ecosystem. Spatial-temporal distribution of coprophagous dipterans communities and their ecosystem functions should be better understood for more efficient Integrated Pest Management (IPM) programs. We also suggest further studies on vectorial capacity and efficiency of certain dung associated Diptera (e.g. *M. ventrosa*) to determine their possibility in transmitting medical and veterinary pathogens.

#### Acknowledgements

We would like to express our deepest gratitude to the management staff and veterinarian of the horse farm who granted permission to us for collecting and studying the dung associated arthropods in their facilities. Thanks to Mr. Haji Sulaiman Abdullah (Faculty of Medicine, UiTM) for his kind assistance in the field work. We would like to thank Dr. Frans Janssens (University of Antwerp, Belgium) for

Collembola identification and Mr. Edward G. Riley (Texas A&M University, USA) for his consultation on beetle identification from pictures. All laboratory facilities provided by the Faculty of Medicine, Universiti Teknologi MARA are greatly acknowledged. Authors also express sincere gratitude to Dr. Meenakshi Bharti and other anonymous reviewer for reviewing the manuscript.

#### References

- Ajwa, H.A. and Tabatabai, M.A. 1994. Decomposition of different organic materials in soils. *Biology and Fertility of Soils* 18(3): 175-182.
- Akhtar, S., Shakeel, S., Mehmood, A., Hamid, A. and Saif, S. 2013. Comparative Analysis of Animal Manure for Soil Conditioning. *International Journal of Agronomy and Plant Production* 4(12): 3360-3366.
- Axtell, R.C. 1963. Acarina occurring in domestic animal manure. *Annals of the Entomological Society of America* 56(5): 628-633.
- Bai, M.G. and Sankaran, T. 1977. Parasites, predators and other arthropods associated with *Musca domestica* and other flies breeding in bovine manure. *Entomophaga* 22(2): 163-167.
- Bay, D.E. and Harris, R.L. 1988. Introduction to veterinary entomology. Bryan, Texas: Stonefly Publishing.p56.
- Bharti, M. 2008. Current status of family Muscidae (Diptera) from India. *Journal of Entomological Research* 32(2): 171-176.
- Blume, R.R. 1985. A checklist, distributional record, and annotated bibliography of the insects associated with bovine droppings on pastures in America north of Mexico. *Southwestern Entomologist* (Supplement 9). Texas A&M University. 55pp.
- Borkent, C.J. and Harder, L.D. 2007. Flies (Diptera) as pollinators of two dioecious plants: behaviour and implications for plant mating. *The Canadian Entomologist* 139(2): 235-246.
- Bøving, A.G. and Henriksen, K.L. 1938. The developmental stages of the Danish Hydrophilidae (Ins. Coleoptera). *Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening i København* 102: 27-162.

- Bowman, D.D. 2014. Georgis' parasitology for veterinarians. 10th Edition. St. Louis, Missouri: Elsevier Health Sciences. pp. 11-80.
- Chaiwong, T., Srivoramas, T., Sueabsamran, P., Sukontason, K., Sanford, M.R. and Sukontason, K.L. 2014. The blow fly, *Chrysomya megacephala*, and the house fly, *Musca domestica*, as mechanical vectors of pathogenic bacteria in Northeast Thailand. *Tropical Biomedicine* 31(2): 336-346.
- Chen, W.Y., Hung, T.H. and Shiao, S.F. 2004. Molecular identification of forensically important blow fly species (Diptera: Calliphoridae) in Taiwan. *Journal of Medical Entomology* 41(1): 47-57.
- Chernova, N.M., Bokova, A.I., Varshav, E.V., Goloshchapova, N.P. and Savenkova, Y.Y. 2007. Zoophagy in Collembola. *Entomological Review* 87: 799-811.
- Chew, W.K., Kurahashi, H., Nazni, W.A., Heo, C.C., Heah, S.K., Jeffery, J. and Lee, H.L. 2012. A new record for *Lispe orientalis* Wiedemann, 1824 (Diptera: Muscidae) from peninsular Malaysia. *Tropical Biomedicine* 29(3): 1-4.
- Cleton, N., Koopmans, M., Reimerink, J., Godeke, G.J. and Reusken, C. 2012. Come fly with me: review of clinically important arboviruses for global travelers. *Journal of Clinical Virology* 55(3): 191-203.
- Costa, M.C. and Weese, J.S. 2012. The equine intestinal microbiome. *Animal Health Research Reviews* 13(1): 121-128.
- Couri, M.S., Pont, A.C. and Penny, N.D. 2006. Muscidae (Diptera) from Madagascar: identification keys, descriptions of new species and new records. *Proceeding of California Academy of Sciences* 57(29): 799-923.
- Dowd, S.E., Callaway, T.R., Wolcott, R.D., Sun, Y., McKeenan, T., Hagevoort, R.G. and Edrington, T.S. 2008. Evaluation of the bacterial diversity in the feces of cattle using 16S rDNA bacterial tag-encoded FLX amplicon pyrosequencing (bTEFAP). *BMC Microbiology* 8(1): 125.
- Ebejer, M.J. and Gatt, P. 1999. The species of Fanniidae and Muscidae of the Maltese Islands. *Studia Dipterologica* 6(1): 79-92.
- Emberson, R.M. (1980). Macrochelidae from the Kermadec Islands and a key to species of *Macrocheles* Latreille from the New Zealand region (Acari: Mesostigmata). *New Zealand Entomologist* 7(2): 135-138.
- Emden, F.I. van. 1965. The fauna of India and the adjacent countries. Diptera, Vol. 7, Muscidae, Part I. Zoological Survey of India. 647 pp.
- Estes, A.M., Hearn, D.J., Snell-Rood, E.C., Feindler, M., Feeser, K., Abebe, T., Dunning Hotopp, J.C. and Moczek, A.P. 2013. Brood ball-mediated transmission of microbiome members in the dung beetle, *Onthophagus taurus* (Coleoptera: Scarabaeidae). *PLoS One* 8(11): e79061.
- Ezenwa, V.O., Gerardo, N.M., Inouye, D.W., Medina, M. and Xavier, J.B. 2012. Animal behavior and the microbiome. *Science* 338(6104): 198-199.
- Fadzil, M. 1973. *Musca conducens* Walker 1859-its prevalence and potential as a vector of *Stephanofilaria kaeli* in peninsular Malaysia. *Kajian Veterinaire* 5: 27-37.
- Finley, S.J., Benbow, M.E. and Javan, G.T. 2015. Potential applications of soil microbial ecology and next-generation sequencing in criminal investigations. *Applied Soil Ecology* 88: 69-78.
- Gestmann, F., Förster, M., Mehlhorn, H., Sievert, K., Messler, S., Neuhausen, N., Petersdorf, S. and Pfeffer, K. 2012. Flies as vectors of microorganisms potentially inducing severe diseases in humans and animals. *In: H. Mehlhorn, (ed.) Arthropods as vectors of emerging diseases.* Springer Berlin Heidelberg. pp. 195-226.
- Getachew, S., Gebre-Michael, T., Erko, B., Balkew, M. and Medhin, G. 2007. Non-biting cyclorrhaphan flies (Diptera) as carriers of intestinal human parasites in slum areas of Addis Ababa, Ethiopia. *Acta Tropica* 103: 186-194.
- Girija, D., Deepa, K., Xavier, F., Antony, I. and Shidhi, P.R. 2013. Analysis of cow dung microbiota-A metagenomic approach. *Indian Journal of Biotechnology* 12(3): 372-378.
- Glida, H. and Bertrand, M. 2002. The occurrence of *Macrocheles* mites (Acari: Macrochelidae) in relation to the activity of dung beetles: a field study in southern France. *In: F. Bernini, R. Nannelli, G. Nuzzaci, and E.D. Lillo, (eds.). Acarid Phylogeny and Evolution: Adaptation in Mites and Ticks.* Springer Netherlands. pp. 199-207.

## Coprophilic dipteran community associated with horse dung in Malaysia

- Graczyk, T., Knight, R., Gilman, R. and Cranfield, M. 2001. The role of non-biting flies in the epidemiology of human infectious diseases. *Microbes and Infection* 3(3): 231-235.
- Greenberg, B. 1973. Flies and disease. Vol. 2. Biology and disease transmission. Princeton University Press. pp 154-339.
- Greenslade, P. and Ireson, J.E. 1986. Collembola of the southern Australian culture steppe and urban environments: a review of their pest status and key to identification. *Journal of the Australian Entomological Society* 25: 273-291.
- Gubler, D.J. 2002. The global emergence/resurgence of arboviral diseases as public health problems. *Archives of Medical Research* 33(4): 330-342.
- Hall, M. and Wall, R. 1995. Myiasis of humans and domestic animals. *Advances in Parasitology* 35: 257-334.
- Hammer, O. 1941. Biological and ecological investigations on flies associated with pasturing cattle and their excrement. *Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening i København* 105: 1-257
- Hanski, I. 1983. Distributional ecology and abundance of dung and carrion-feeding beetles (Scarabaeidae) in tropical rain forests in Sarawak, Borneo. *Acta Zoologica Fennica* 167: 1-45.
- Hanski, I. and Cambefort, Y. 1991. Dung beetle ecology. Princeton University Press. pp. 3-68.
- Hanski, I. and Koskela, H. 1979. Resource partitioning in six guilds of dung-inhabiting beetles (Coleoptera). *Annales Entomologici Fennici* 45: 1-12.
- Heo, C.C., Nazni, W.A., Chew, W.K., Kurahashi, H., Jeffery, J., Heah, S.K., and Omar, B. 2010. A Study of Cow Dung Diptera in Sentul Timur, Kuala Lumpur, Malaysia. *The Journal of Tropical Medicine and Parasitology* 33(2): 53-61.
- Hirabayashi, K. and Wotton, R.S. 1998. Organic matter processing by chironomid larvae (Diptera: Chironomidae). *Hydrobiologia* 382(1-3): 151-159.
- Holter, P. 1977. An experiment on dung removal by *Aphodius* larvae (Scarabaeidae) and earthworms. *Oikos* 28: 130-136.
- Holter, P. 1979. Effect of dung-beetles (*Aphodius* spp.) and earthworms on the disappearance of cattle dung. *Oikos* 32: 393-402.
- Howard, J. 2001. Nuisance flies around a landfill: patterns of abundance and distribution. *Waste Management and Research* 19(4): 308-313.
- Hulley, P.E. 1986. Factors affecting numbers of *Musca domestica* Linnaeus (Diptera: Muscidae) and some other flies breeding in poultry manure. *Journal of Entomological Society of South Africa* 49(1): 19-27.
- Hussey, N.W. 1957. Observations on the Sphaeroceridae (Borboridae, Diptera) associated with stable manure. *Journal of Natural History* 10(118): 762-766.
- James, M.T. 1947. The flies that cause myiasis in man. US Department of Agriculture. Misc. Pub. No. 631. 175 pp.
- Johnson, S.J., Parker, R.J., Norton, J.H., Jaques, P.A. and Grimshaw, A.A. 1981. Stephanofilariasis in cattle. *Australian Veterinary Journal* 57(9): 411-413.
- Khoso, F.N., Wong, S.K., Chia, S.L. and Lau, W.H. 2015. Assessment of non-biting synanthropic flies associated with fresh markets. *Journal of Entomology and Zoological Studies* 3(1): 13-20.
- Koehler, H.H. 1999. Predatory mites (Gamasina, Mesostigmata). *Agriculture, Ecosystems and Environment* 74(1): 395-410.
- Krantz, G.W. 1983. Mites as biological control agents of dung-breeding flies, with special reference to the Macrochelidae. In: Hoy M.A. (ed.), *Biological Control of Pests by Mites*. University of California, Berkeley. pp. 91-98.
- Krantz, G.W. and Walter, D.E. 2009. *A manual of acarology*. Lubbock: Texas Tech University Press. 807 pp.
- Kumara, T.K., Disney, R.H.L., Hassan, A.A., Flores, M., Hwa, T.S., Mohamed, Z., CheSalmah, M.R. and Bhupinder, S. 2012. Occurrence of oriental flies associated with indoor and outdoor human remains in the tropical climate of north Malaysia. *Journal of Vector Ecology* 37(1): 62-68.
- Kurahashi, H. and Shinonaga, S. 2009. Two new species of the muscid flies from Sarawak, East Malaysia (Diptera: Muscidae). *Japanese Journal of Systematic Entomology* 15(2): 299-306.



- Laurence, B.R. 1955. The ecology of some British Sphaeroceridae (Borboridae, Diptera). *The Journal of Animal Ecology* 24(1): 187-199.
- Lee, H.L., Krishnasamy, M., Abdullah, A.G. and Jeffery, J. 2004. Review of forensically important entomological specimens in the period of 1972–2002. *Tropical Biomedicine* 21(2): 69-75.
- Mašán, P. and Halliday, B. 2009. Mesostigmatid mites associated with the dung beetle *Copris lunaris* (Coleoptera: Scarabaeidae). *European Journal of Entomology* 106(4): 545-550.
- Nazni, W.A., Apandi, M.Y., Eugene, M., Azahari, A.H., Shahar, M.K., Zainah, S., Vthylingam, I. and Lee, H.L. 2013. The potential of house fly, *Musca domestica* (L.) in the mechanical transmission of influenza A subtype H1N1 virus under laboratory conditions. *Journal of General and Molecular Virology* 5(2): 22-28.
- Nazni, W.A., Nooraidah, H., Jeffery, J., Azahari, A.H., Mohd Noor, I., Sadiyah, I. and Lee, H.L. 2007. Distribution and abundance of diurnal and nocturnal dipterous flies in the Federal Territory, Putrajaya. *Tropical Biomedicine* 24: 61-66.
- Nazni, W.A., Seleena, B., Lee, H.L., Jeffery, J., Rogayah, T.A.R. and Sofian, M.A. 2005. Bacteria fauna from the house fly, *Musca domestica* (L.). *Tropical Biomedicine* 22(2): 225-231.
- Nichols, E., Larsen, T., Spector, S., Davis, A.L., Escobar, F., Favila, M., Vulinec, K. and Network, T.S.R. 2007. Global dung beetle response to tropical forest modification and fragmentation: a quantitative literature review and meta-analysis. *Biological Conservation* 137(1): 1-19.
- Nichols, E., Spector, S., Louzada, J., Larsen, T., Amezquita, S., Favila, M.E. and Network, T.S.R. 2008. Ecological functions and ecosystem services provided by Scarabaeinae dung beetles. *Biological Conservation* 141(6): 1461-1474.
- Nurita, A.T., and Hassan, A.A. 2013. Filth flies associated with municipal solid waste and impact of delay in cover soil application on adult filth fly emergence in a sanitary landfill in Pulau Pinang, Malaysia. *Bulletin of Entomological Research* 103(3): 296-302.
- Papp, L. 2008. A review of the Old World *Coproica* Rondani, 1861 (Diptera, Sphaeroceridae), with descriptions of twelve new species. *Acta Zoologica Academiae Scientiarum Hungaricae* 54(2): 47-209.
- Patnaik, B. and Roy, S.P. 1969. Studies on stephanofilariasis in Orissa. IV. rearing of *Musca conducens* Walker, 1859, the arthropod vector of *S. assamensis* in the laboratory. *Indian Journal of Animal Sciences* 39(6): 529-535.
- Patton, W.S. 1922. Notes on the species of the genus *Musca* Linnaeus. Part 1. *Bulletin of Entomological Research* 12:411-426.
- Pechal, J.L., Crippen, T.L., Benbow, M.E., Tarone, A.M., Dowd, S. and Tomberlin, J.K. 2014. The potential use of bacterial community succession in forensics as described by high throughput metagenomic sequencing. *International Journal of Legal Medicine* 128(1): 193-205.
- Pont, A.C., Harutyunova, K., Harutyunova, M. and Werner, D. 2012. The hunter-flies of Armenia. II. The genus *Lispe* Latreille, 1796: (Insecta: Diptera: Muscidae). *Zoology in the Middle East* 55(1): 79-84.
- Putman, R.J. 1978. The role of carrion-frequenting arthropods in the decay process. *Ecological Entomology* 3(2): 133-139.
- Reid, J.A. 1953. Notes on house-flies and blow-flies in Malaya. *Bulletin of Institute for Medical Research Malaya* 7: 1-26.
- Service, M. 2012. *Medical entomology for students*. 5<sup>th</sup> Edition. Cambridge: Cambridge University Press. 317 pp.
- Shade, A., Mc Manus, P.S. and Handelsman, J. 2013. Unexpected diversity during community succession in the apple flower microbiome. *MBio* 4(2): e00602-12.
- Shinonaga, S. and Kano, R. 1989. Four new species of *Lispe* (Diptera: Muscidae) from the Oriental Region. *Japanese Journal of Entomology* 57(4): 815-821.
- Shinonaga, S. and Pont, A.C. 1992. A new species of the genus *Lispe* Latreille, with notes on two related species, *L. assimilis* Wiedemann and *L. microptera* Seguy (Diptera: Muscidae). *Japanese Journal of Entomology* 60(4): 715-722.
- Sommer, C., Jensen, K.M.V. and Jespersen, J.B. 2001. Topical treatment of calves with



## Coprophilic dipteran community associated with horse dung in Malaysia

- synthetic pyrethroids: effects on the non-target dung fly *Neomyia cornicina* (Diptera: Muscidae). Bulletin of Entomological Research 91(2): 131-137.
- Spencer, K.A. 1973. Agromyzidae (Diptera) of economic importance. Vol. 9. Springer Science and Business Media. pp. 11-28.
- Stoffolano, J.G. and Geden, C.J. 1987. Succession of manure arthropods at a poultry farm in Massachusetts, USA, with observations on *Carcinops pumilio* (Coleoptera: Histeridae) sex ratios, ovarian condition, and body size. Journal of Medical Entomology 24(2): 212-220.
- Sucharit, S. and Tumrasvin, W. 1981. The survey of flies of medical and veterinary importance in Thailand. Japanese Journal of Sanitary Zoology 32(4): 281-285.
- Sucharit, S., Kerdpibule, V., Tumrasvin, W., Deesin, T. and Nakorn, C.N. 1981. Myiasis of the vagina of a comatose woman caused by *Parasarcophaga ruficornis* Fabricius. Journal of the Medical Association of Thailand 64(11): 580-583.
- Sukontason, K.L., Narongchai, P., Lerthamngtham, S., Piangjai, S. and Olson, J.K. 2002. *Chrysomya rufifacies* (Macquart) as a forensically-important fly species in Thailand: a case report. Journal of Vector Ecology 26(2): 162-164.
- Sukontason, K.L., Narongchai, P., Sripakdee, D., Boonchu, N., Chaiwong, T., Ngern-Klun, R., Piangjai, S. and Sukontason, K. 2005. First report of human myiasis caused by *Chrysomya megacephala* and *Chrysomya rufifacies* (Diptera: Calliphoridae) in Thailand, and its implication in forensic entomology. Journal of Medical Entomology 42(4): 702-704.
- Sukontason, K., Narongchai, P., Kanchai, C., Vichairat, K., Sribanditmongkol, P., Bhoopat, T., Kurahashi, H., Chockjamsi, M., Piangjai, S., Bunchu, N., Vongvivach, S., Samai, W., Chaiwong, T., Methanitikorn, R., Ngern-Klun, R., Sripakdee D., Boonsriwong W., Siriwananarungsee S, Srimuangwong, C., Hanterdsith, B., Chaiwan, K., Srisuwan, C., Upakut, S., Moopayak, K., Vogtsberger, R.C., Olson, J.K., Sukontason, K.L. 2007. Forensic entomology cases in Thailand: a review of cases from 2000 to 2006. Parasitology Research 101(5): 1417-1423.
- Sulaiman, S., Othman, M.Z. and Aziz, A.H. 2000. Isolations of enteric pathogens from synanthropic flies trapped in downtown Kuala Lumpur. Journal of Vector Ecology 25(1): 90-93.
- Sulaiman, S., Sohadi, A.R., Yunus, H. and Iberahim, R. 1988. The role of some cyclorrhaphan flies as carriers of human helminths in Malaysia. Medical and Veterinary Entomology 2(1): 1-6.
- Tan, S.H., Rizman-Idid, M., Mohd-Aris, E., Kurahashi, H. and Mohamed, Z. 2010. DNA-based characterisation and classification of forensically important flesh flies (Diptera: Sarcophagidae) in Malaysia. Forensic Science International 199(1): 43-49.
- Taylor, D.B., Friesen, K., Zhu, J.J. and Sievert, K. 2012a. Efficacy of cyromazine to control immature stable flies (Diptera: Muscidae) developing in winter hay feeding sites. Journal of Economic Entomology 105(2): 726-731.
- Taylor, D.B., Moon, R.D. and Mark, D.R. 2012b. Economic impact of stable flies (Diptera: Muscidae) on dairy and beef cattle production. Journal of Medical Entomology 49(1): 198-209.
- Tischler W. 1949: Grundzüge der terrestrischen Tierökologie. Braunschweig: Friedrich Vieweg und Sohn. (In German). 219 pp.
- Tomberlin, J.K., Crippen, T.L., Tarone, A.M., Singh, B., Adams, K., Rezenom, Y.H., Benbow, M. E., Flores, M., Longnecker, M., Pechal, J.L., Russell, D.H., Beier, R.C. and Wood, T.K. 2012. Interkingdom responses of flies to bacteria mediated by fly physiology and bacterial quorum sensing. Animal Behaviour 84(6): 1449-1456.
- Triplehorn, C.A. and Johnson, N.F. 2005. Borror and DeLong's Introduction to the Study of Insects. Belmont, CA: Thomson Brooks/Cole. 864 pp.
- Tumrasvin, W. and Shinonaga, S. 1982. Studies on medically important flies in Thailand VII Report on 73 species of muscid flies (excluding Muscinae and Stomoxyinae) with the taxonomic keys (Diptera: Muscidae). Japanese Journal of Sanitary Zoology 33: 181-199.

- Valiela, I. 1969. An experimental study of the mortality factors of larval *Musca autumnalis*. De Geer Ecological Monograph 39: 199-220.
- Valiela, I. 1974. Composition, food webs and population limitation in dung arthropod communities during invasion and succession. American Midland Naturalist 92(2): 370-385.
- Vikhrev, N.E. 2012a. Four new species of *Lispe* Latreille, 1976 (Diptera: Muscidae) with taxonomic notes on related species. Russian Entomological Journal 21(4): 423-433.
- Vikhrev, N.E. 2012b. Revision of the *Lispe longicollis*-group (Diptera, Muscidae). ZooKeys 235: 23-29.
- Wang, J., Li, Z., Chen, Y., Chen, Q. and Yin, X. 2008. The succession and development of insects on pig carcasses and their significances in estimating PMI in south China. Forensic Science International 179(1): 11-18.
- Wang, Y.X. and Feng, Y. 2008. Investigation on the calypterae flies in Mountain Mengdingshan, Mingshan County, Sichuan Province, China. Entomological Journal of East China 3: 236-240.
- Watrelet-Virieux, D. and Pin, D. 2006. Chronic eosinophilic dermatitis in the scrotal area associated with stephanofilariasis infestation of charolais bull in France. Journal of Veterinary Medicine 53(3): 150-152.
- Werner, D. and Pont, A. 2006. New results on Diptera predator in the blackfly plaque areas of central Europe and the caucasus. Acta Entomologica Serbica 4: 131-140.
- White, I.M. and Elson-Harris, M.M. 1992. Fruit flies of economic significance: their identification and bionomics. CAB International. 601 pp.
- Yap, K.L., Kalpana, M. and Lee, H.L. 2008. Wings of the common house fly (*Musca domestica* L.): importance in mechanical transmission of *Vibrio cholerae*. Tropical Biomedicine 25(1): 1-8.
- Yoshihara, Y. and Sato, S. 2015. The relationship between dung beetle species richness and ecosystem functioning. Applied Soil Ecology 88: 21-25.
- Zhang, J.J., He, M.X., Jia, S.B. and Wang, L.Y. 2012. Spatial distribution and biological effects of springtail *Ceratophysella duplicispinosa* in Chinese cabbage plot. Northern Horticulture 4: 1-3.